



ARTICLE

Anti-Biofilm Effect of Egg Yolk Phosvitin by Inhibition of Biomass Production and Adherence Activity against *Streptococcus mutans*

Hyeon Joong Kim¹, Jae Hoon Lee¹, Dong Uk Ahn², and Hyun-Dong Paik^{1,*}

¹Department of Food Science and Biotechnology of Animal Resources, Konkuk University, Seoul 05029, Korea

²Department of Animal Science, Iowa State University, Ames, IA 50011, USA

OPEN ACCESS

Received July 31, 2020

Revised August 12, 2020

Accepted August 18, 2020

*Corresponding author : Hyun-Dong Paik
Department of Food Science and
Biotechnology of Animal Resources,
Konkuk University, Seoul 05029, Korea
Tel: +82-2-2049-6011
Fax: +82-2-455-3082
E-mail: hdpai@konkuk.ac.kr

*ORCID
Hyeon Joong Kim
<https://orcid.org/0000-0001-6999-868X>
Jae Hoon Lee
<https://orcid.org/0000-0002-7440-6842>
Dong Uk Ahn
<https://orcid.org/0000-0001-9523-7630>
Hyun-Dong Paik
<https://orcid.org/0000-0001-9891-7703>

Abstract The formation of biofilms on the enamel surface of teeth by *Streptococcus mutans* is an important step in dental plaque formation, demineralization, and early caries because the biofilm is where other bacteria involved in dental caries attach, grow, and proliferate. The objectives of this study were to determine the effect of phosvitin (PSV) on the biofilm formation, exopolysaccharides (EPS) production, adherence activity of *S. mutans*, and the expression of genes related to the compounds essential for biofilm formation (quorum-sensing inducers and components of biofilm matrix) by *S. mutans*. PSV significantly reduced the biofilm-forming activity of *S. mutans* and increased the degradation of preformed biofilms by *S. mutans*. PSV inhibited the adherence activity of *S. mutans* by 31.9%–33.6%, and the production of EPS by 62%–65% depending upon the strains and the amount of PSV added. The expressions of genes regulating the production of EPS and the quorum-sensing-inducers (*gtfA*, *gtfD*, *ftf*, *relA*, *vicR*, *brpA*, and *comDE*) in all *S. mutans* strains were down-regulated by PSV, but *gtfB* was down-regulated only in *S. mutans* KCTC 5316. Therefore, the anti-biofilm-forming activity of PSV was accomplished through the inhibition of biofilm formation, adherence activity, and the production of quorum-sensing inducers and EPS by *S. mutans*.

Keywords phosvitin, *Streptococcus mutans*, biofilm formation

Introduction

The initial step of dental caries is the colonization of *Streptococcus mutans*, a gram-positive, facultative anaerobic bacteria, on the surface of human teeth and forming biofilms (Banas, 2004; Burne, 1998; Smith et al., 1993) because the biofilm is where other bacteria involved in dental caries attach, grow, and proliferate (Hamada and Slade, 1980). Also, the biofilm protects the bacteria from environmental stresses, antibacterial agents, and host antibodies (Stewart, 1996). The production of exopolysaccharides (EPS) and acids from carbohydrates is another important step for the cariogenic process by *S. mutans* (Koo et al., 2003). *S. mutans* use fructosyltransferase (FTF) (Shemesh and

Steinberg, 2006) and glucosyltransferases (GTFs) to produce fructans and glucans from sucrose (Bowen and Koo, 2011). Fructans are extracellular storage compounds that act as binding sites for bacteria (Burne et al., 1996). Glucans play key roles in the attachment and colonization of *S. mutans* on the surface of the tooth, which is essential for the structural establishment of biofilms (Koo et al., 2003). Bacteria use a method called quorum-sensing to control their metabolic activity. Gene expression and communications among bacteria are important tools for them to adapt to various environmental conditions and defend themselves from the competitors and host antibodies (Shapiro, 1998). The expression of genes that regulate the production of exopolysaccharides (fructans and glucans), components of biofilm matrix, and quorum-sensing inducers in *Streptococcus* affects biofilm formation and dental diseases by regulating several physiological properties (Al-Sohaibani and Murugan, 2012). To prevent dental caries, therefore, both inhibiting biofilm formation by *S. mutans* and controlling dietary factors are important (Bradshaw and Lynch, 2013).

Most of the previous research efforts on dental caries were focused on controlling the fermentable dietary carbohydrates that serve as substrates for dental plaque (Paes Leme et al., 2006). However, foods also contain a variety of bioactive compounds that provide biological benefits (Siriwardhana et al., 2013): milk, tea, apples, and algae are considered to have beneficial effects on teeth, and some food-derived ingredients are known to inhibit caries through anti-bacterial and anti-biofilm activities (Daglia et al., 2010; Gazzani et al., 2012; Rajaraman et al., 2014; Taborsky and Mok, 1967). The egg is a well-known source of minerals, lipids, proteins, and biologically active peptides (Mine, 2007), and some of the egg proteins were found effective in inhibiting biofilm formation by preventing the adherence of various microorganisms, EPS production, demineralization, pH reduction, and biomass formation (Bradshaw and Lynch, 2013, Giacaman et al., 2019).

Phosvitin (PSV) is the major phosphoglycoprotein in egg yolk (Abe et al., 1982) with a unique structural characteristic: it contains a large number of phosphoserine groups (55% of the total amino acids) in its structure. Thus, PSV has a very strong metal-binding capability (Samaraweera et al., 2011), and shows strong antimicrobial (Lee et al., 2002) and antioxidant activities (Choi et al., 2004). The objectives of this study were to determine the effects of PSV on 1) the biofilm formation, 2) the removal of preformed biofilm, 3) the adherence ability, and 4) the expression of genes regulating the production of quorum-sensing inducers and the matrix-forming compounds (EPS) by *S. mutans*.

Materials and Methods

Materials

PSV was isolated from chicken egg yolk following the method of Lee et al. (2014). Crystal violet was purchased from Alfa Aesar (Haverhill, MA, USA), brain heart infusion broth (BHI) from HiMedia Laboratories (Mumbai, India), Congo red from Sigma-Aldrich (St. Louis, MO, USA), and SYBR green from Bioline (London, UK), TRIzolTM MaxTM Bacterial RNA Isolation Kit from Invitrogen (Carlsbad, CA, USA), Revert Aid First-strand cDNA synthesis Kit from Thermo Fisher Scientific (Waltham, MA, USA). The *S. mutans* KCTC 5124, 5458, and 5316 strains were used in this study. All the strains were grown in BHI media at 37°C. The cultures were stored at -80°C in BHI containing 25% glycerol.

Biofilm formation

Biofilm formation was estimated using the crystal violet assay in a 96-well microplate (Kulshrestha et al., 2016). *S. mutans* were cultivated overnight in BHI media. *S. mutans* were diluted to 10⁵-10⁶ colony forming units (CFU)/mL. Fresh media (BHI+0.1% sucrose) containing 0, 0.25, 0.5, and 1 mg/mL of PSV (100 µL) and diluted *S. mutans* (100 µL) were

added into microplate wells and the samples were incubated at 37°C under anaerobic conditions for 24 h. After incubation, planktonic cells and media were removed by gently rinsing with sterile distilled water three times. The adhered biofilm was fixed by adding 100 µL methanol for 15 min. After fixing, each well was rinsed with sterile distilled water three times and stained with 100 µL of 0.1% crystal violet for 5 min at room temperature. After staining, crystal violet was removed gently and rinsed with sterile distilled water three times. Dimethyl sulfoxide (100 µL) was added to the wells to dissolve the stained biofilm, and the absorbance was determined at 570 nm using a microplate reader (model 680, BioRad, Hercules, CA, USA).

Preformed biofilm

Overnight cultured *S. mutans* diluted to 10⁵–10⁶ CFU/mL in fresh BHI media (0.1% sucrose) and 200 µL of *S. mutans* were inoculated to each well of a 96-well microplate. The samples were incubated at 37°C under anaerobic conditions for 24 h. The planktonic cells and media were removed and rinsed with phosphate-buffered saline (PBS). PSV at 0, 0.25, 0.5, and 1 mg/mL and media (BHI+0.1% sucrose) were added to each well and the microplate and the samples were incubated at 37°C under anaerobic conditions for 24 h. After incubation, the effects of PSV on biofilm was estimated by the crystal violet assay.

Bacterial adherence

The glass surface adherence assay was used to evaluate the effects of PSV on bacterial adherence (Hamada and Slade, 1980). *S. mutans* 10⁵–10⁶ CFU/mL were transferred to glass tubes containing BHI media (0.1% sucrose) and PSV. The tubes were tilted 30 degrees and then incubated at 37°C under anaerobic conditions for 24 h. After incubation, the attached bacteria were diluted with 0.5 M sodium hydroxide and the supernatant containing planktonic cells was collected. The adhered cells and total cells were estimated by reading the absorbance at 600 nm using a spectrophotometer (Thermo Fisher Scientific). The bacterial adherence was calculated using the following formula:

$$\text{Bacterial adherence (\%)} = (\text{OD of adherent cells} / \text{OD of total cells}) \times 100$$

Bacterial exopolysaccharide production

Bacterial exopolysaccharide (EPS) production was evaluated by the Congo red (CR)-binding assay (Smalley et al., 1995). *S. mutans* 10⁵–10⁶ CFU/mL, media (BHI+0.1% sucrose, 200 µL) containing 0, 0.25, 0.5, and 1 mg/mL of PSV and Congo red dye (0.5 mM, 50 µL) were added to each well of a microplate, and incubated at 37°C for 1 h. At the end of incubation, the supernatant in each well was transferred to individual microtubes and centrifuged at 10,000 g for 5 min. The supernatant (200 µL) of each tube was collected and the absorbance read at 490 nm. Bacterial EPS produced was estimated using the following formula:

$$\text{EPS production (\%)} = (\text{OD of blank CR} - \text{OD of the supernatant}) / \text{OD of the control} \times 100$$

Quantitative real-time PCR analysis

Quantitative real-time polymerase chain reaction (qRT-PCR) was performed to evaluate the effect of PSV on the

expression of the genes that regulate the production of biofilm matrix components and quorum sensing inducers in *S. mutans*. Total RNA was isolated from *S. mutans* strains using RNA Isolation Kit and the RNA concentration was estimated at 260 nm by a spectrophotometer. cDNA was synthesized using the cDNA Synthesis Kit. The virulence gene primer sequences are shown in Table 1 (Koolheat et al., 2016) and 16S rRNA primer used for reference gene. The SYBR green reagent was used to determine the amount of DNA. The PCR was performed for 40 cycles with a denaturing temperature at 95°C for 5 s and annealing/extension temperature at 65°C for 30 s. The qRT-PCR results were analyzed using the qRT-PCR software (PikoReal software 2.2, Thermo Fisher Scientific) that evaluates the expression of the virulence gene. The purity of PCR products was estimated using the melting curve.

Scanning electron microscopy

Overnight cultured *S. mutans* (10^5 – 10^6 CFU/mL), media (BHI+0.1% sucrose, 200 µL) and PSV (1 mg/mL, 200 µL) were incubated at 37°C for 24 h in a 6-well microplate with glass coverslips. After incubation, the supernatants in the wells were removed and the biofilm on each glass coverslip was washed with PBS. The biofilms on the glass coverslips were fixed with 2.5% glutaraldehyde at 4°C for 1 h, washed three times with PBS and dehydrated in various concentrations of ethanol (from 50% to 100%) for 15 min. The dehydrated coverslips were dipped in isoamyl acetate for 15 min and then freeze-dried. The biofilm was scanned using a field emission scanning electron microscope (FESEM) (SU8010, Hitachi High-Technologies, Tokyo, Japan).

Statistical analysis

All results were presented as the means and the standard deviation of three replicates. The results of biofilm formation, preformed biofilm formation, EPS production, and adherence activity were performed one-way analysis of variance (ANOVA) to measure the significance of the difference in means. The student's t-test was used to measure the significance of the difference in the means of qRT-PCR results. All results were calculated using IBM SPSS Statistics (Version 25, IBM Corp, Armonk, NY, USA).

Table 1. Primer sequences used in quantitative real-time PCR

Gene	Description	Forward primer (5' → 3')	Reverse primer (5' → 3')
<i>gtfA</i>	Sucrose phosphorylase	AGGAAGTGAAGCGGCCAGT	TCAATACGGCCATCCAAATCA
<i>gtfB</i>	Glucosyltransferase B	AGCAATGCAGCCAATCTACAAAT	ACGAACTTGCCGTTATTGTCA
<i>gtfD</i>	Glucosyltransferase-I	ACAGCAGACAGCAGCCAAGA	ACTGGGTTGCTGCGTTG
<i>ftf</i>	Fructosyltransferase	AAATATGAAGGCGGCTACAACG	CTTCACCAGTCTTAGCATCCTGAA
<i>relA</i>	Guanosine tetra (penta)-phosphatesynthetase	ACAAAAAGGGTATCGTCCGTACAT	AATCACCGCTTGGTATTGCTAATTG
<i>vicR</i>	Histidine kinase two-component regulatory system	TGACACGATTACAGCCTTGATG	CGTCTAGTTCTGGTAACATTAAGTCC AATA
<i>brpA</i>	Biofilm-regulation protein	GGAGGAGCTGCATCAGGATTC	AACTCCAGCACATCCAGCAAG
<i>comDE</i>	Competence-stimulating system	ACAATTCCCTTGAGTTCCATCCAAG	TGGTCTGCTGCCTGTTGC
16S rRNA	16S ribosomal RNA, normalizing internal standard	CCTACGGGAGGCAGCAGTAG	CAACAGAGCTTACGATCCGAAA

Results

Inhibition of biofilm formation and the degradation of preformed biofilm by PSV

The biofilm formation and maintenance properties are an important factor of surviving in *S. mutans*. The inhibitory activity of PSV on the formation of biofilm by *S. mutans* is shown in Fig. 1. The addition of PSV at 0.25, 0.5, and 1.0 mg/mL inhibited the biofilm formation of *S. mutans* KCTC 5124 strain by 34.0%, 35.7%, and 42.5%, respectively, compared with the control. With the *S. mutans* KCTC 5458 strain, the reductions were 43.9%, 53.5%, and 58.8%, respectively, and 25.4%, 30.1%, and 33.2% with *S. mutans* KCTC 5316 strain. As the concentration of PSV increased, the biofilm formation by *S. mutans* gradually decreased, suggesting a dose-dependent anti-biofilm forming activity of PSV in *S. mutans* ($p<0.05$). The amount of preformed biofilm by *S. mutans* KCTC 5124 strain decreased by 25.6%, 26.9%, and 31.9% compared with the control when 0.25, 0.5, and 1.0 mg/mL of PSV was added (Fig. 2). The reduction of preformed biofilm by *S. mutans* KCTC 5458 strain was 21.5%, 27.5%, and 33.7% when 0.25, 0.5, and 1.0 mg/mL of PSV, respectively, was added, and that by *S. mutans* KCTC 5316 strain was 23.0%, 25.5%, and 33.5%, respectively. These results showed that the PSV inhibited biofilm formation and preformed biofilm of *S. mutans*.

Effect of PSV on the adherence of *S. mutans* on the glass surfaces

The attachment of *S. mutans* on the host tooth surface is first stage and key factor in colonization and biofilm formation. The effect of PSV on the adherence of *S. mutans* performed the glass surfaces adherence assay. Addition of 0.25, 0.5, and 1.0 mg/mL of PSV to the sample reduced 25.5%, 26.9%, and 31.9% of the adherence capacity of *S. mutans* KCTC 5124 strain on the glass surfaces (Fig. 3). For *S. mutans* KCTC 5458 strain, the reduction was 21.5%, 27.5%, and 33.6%, respectively, while that of *S. mutans* KCTC 5316 strain was 23.0%, 25.5%, and 33.5%. The reduction of adherence to the glass surface was dose-




Fig. 1. Effects of phosvitin on biofilm formation against *Streptococcus mutans*. ■, *S. mutans* KCTC 5124, ▨, *S. mutans* KCTC 5458, and ▨, *S. mutans* KCTC 5316. Values are expressed as the mean \pm SD. ^{a-g} Different letters among samples indicate significant differences ($p<0.05$). Control, non-treated group.




Fig. 2. Effects of phosvitin on preformed biofilm against *Streptococcus mutans*. ■, *S. mutans* KCTC 5124, ▨, *S. mutans* KCTC 5458, and ▨, *S. mutans* KCTC 5316. Values are expressed as the mean \pm SD. ^{a-e} Different letters among samples indicate significant differences ($p<0.05$). Control, non-treated group.




Fig. 3. Effects of phosvitin on adherence against *Streptococcus mutans*. ■, *S. mutans* KCTC 5124, ▨, *S. mutans* KCTC 5458, and ▨, *S. mutans* KCTC 5316. Values are expressed as the mean \pm SD. ^{a-g} Different letters among samples indicate significant differences ($p<0.05$). Control, non-treated group.

dependently ($p<0.05$). These results indicated that the PSV inhibited adherence property of *S. mutans*.

Effect of PSV on EPS production

In the presence of PSV, the EPS production by *S. mutans* decreased dose-dependently (Fig. 4) ($p<0.05$). In *S. mutans*




Fig. 4. Effects of phosvitin on EPS production against *Streptococcus mutans*. ■, *S. mutans* KCTC 5124, ▨, *S. mutans* KCTC 5458, and ▨, *S. mutans* KCTC 5316. Values are expressed as the mean±SD. ^{a–e} Different letters among samples indicate significant differences ($p<0.05$). Control, non-treated group; EPS, extracellular polysaccharides.

KCTC 5124 strain, the addition of PSV at 0.25, 0.5, and 1.0 mg/mL reduced the production of EPS by 40.8%, 51.5%, and 64.8%, respectively, compared with the control. The reductions of EPS production by the same PSV treatments to *S. mutans* KCTC 5458 and 5316 strain were 44.4%, 53.7%, and 65.3% and 34.0%, 43.4%, and 62.0%, respectively.

Expression of virulence genes

The effect of PSV on the expression of virulence genes (*gtfA*, *gtfB*, *gtfD*, *ftf*, *vicR*, *relA*, *brpA*, and *comDE*) associated with the biofilm-forming activity of *S. mutans* indicated that all the genes, except for *gtfB*, tested were significantly down-regulated by PSV (Fig. 5). The expression of *gtfB* decreased by 92% in KCTC 5316 strain, but there was no difference between the control group and the PSV treated group in the KCTC 5124 and 5458 strains, respectively. The expression of other virulence genes (*gtfA*, *gtfD*, *ftf*, *relA*, *vicR*, *brpA*, and *comDE*) was significantly down-regulated in all strains of *S. mutans* in the presence of PSV. However, the degree of down-regulation in KCTC 5316 strain was greater than that in other *S. mutans* strains.

Visualization of changes in biofilm through SEM

The scanning electron microscopic (SEM) images of *S. mutans* strains showed dramatic reductions in biofilm formation on the glass surfaces in the presence of 1 mg/mL PSV (Fig. 6). The SEM image of *S. mutans* strains also showed significant reductions in the amounts of biofilms and the numbers of bacteria on the glass surfaces, which agree to the other results in this study - biofilm formation (Fig. 1), bacterial adhesion (Fig. 3) and EPS production (Fig. 4). The SEM image of control (untreated with PSV) showed a biofilm composed of multiple layers with a large number of attached bacteria, while those treated with PSV showed significantly reduced number of the attached bacteria and thin biofilm layers.




Fig. 5. Expression of virulence genes of *Streptococcus mutans* by qRT-PCR. (A) *S. mutans* KCTC 5124, (B) *S. mutans* KCTC 5458 and (C) *S. mutans* KCTC 5316. Values are expressed as the mean \pm SD and analyzed by student's t-test, with $p<0.05$ considered as statistically significant. Control, non-treated group.

Discussion

The biofilm formation by *S. mutans* is the major virulence factor in developing dental caries (Hamada and Slade, 1980). The biofilm formed by *S. mutans* is difficult to remove because it is surrounded by a matrix composed of polysaccharides and other bacterial biofilms. The key to the inhibition of biofilm is suppressing its formation at an early stage (Islam et al., 2008). PSV showed dose-dependent activity in the reduction of the preformed biofilm by *S. mutans* ($p<0.05$). PSV significantly inhibited the formation of biofilm by *S. mutans* at an early stage. Also, the amounts of preformed biofilms by *S. mutans* decreased when PSV was present, suggesting that PSV not only prevented the formation of biofilms but also degraded the preformed biofilms by *S. mutans*. Thus, PSV has great potential to be used to prevent dental caries.

In general, bacteria are preferentially colonized on the surfaces that are hydrophobic and with roughness surfaces (Bowen and Koo, 2011). *S. mutans* starts to colonize on the teeth surface after adhesion using sucrose-dependent and sucrose-independent processes. The sucrose-dependent attachment process is associated with the production of glucans and fructans, while the sucrose-independent attachment process is associated with the electrostatic forces, cell-surface-binding proteins,




Fig. 6. Scanning electron micrographs of *Streptococcus mutans* biofilms on glass coverslips treatment with phosvitin 1 mg/mL. (A) *S. mutans* KCTC 5124, (B) *S. mutans* KCTC 5458 and (C) *S. mutans* KCTC 5316. Control, non-treated group. $\times 10,000$ magnification.

and hydrophobic interactions (Staat et al., 1980). PSV inhibited the adhesion of *S. mutans* cells on the glass surfaces by changing the biofilm matrix. PSV inhibited the sucrose-dependent attachment by down-regulating carbohydrate metabolism and the sucrose-dependent EPS production. Also, PSV inhibited the initial sucrose-independent attachment by down-regulated *comDE* gene and inhibited LytR. The decrease of adhesion is expected to have a marked inhibitory effect on early colonization and biofilm formation by *S. mutans*.

Polysaccharides are the major part of the biofilm matrix and the exopolysaccharides produced by *S. mutans* are the key factor to form indispensable mature biofilms, the maintenance, and expansion of the biofilms (Flemming and Wingender, 2010). Thus, the amounts and types of exopolysaccharides produced are very important for the formation, maintenance of the bacterial community, and the structure of biofilms (Sutherland, 2001). This indicated that the significant reduction of EPS production by *S. mutans* in the presence of PSV is very important for the anti-biofilm-forming ability of PSV.

The downregulation of virulence genes in *S. mutans* by PSV indicated that the inhibitory activity of PSV in the biofilm formation by *S. mutans* is related to its effect on gene expression. The *gtfA* genes encode GTF-A enzyme, a sucrose phosphorylase (Russell et al., 1988). The *gtfB* encodes water-insoluble glucans, the adhesion molecules that are essential for the sucrose-dependent attachment and immobilization of bacteria on hard surfaces (Wen et al., 2010). The *gtfD* produces water-soluble glucans that are sucrose-dependent (Hanada and Kuramitsu, 1988). Although, in KCTC 5124 and 5458 strains PSV did not affect to expression of *gtfB*, gene expression of *gtfA* and *gtfD* was decreased, and similarly, EPS formation was also decreased. Therefore, PSV suppressed the expression and activity of GTFs, which lowered the formation of EPS. These

results indicated that PSV has anti-biofilm activity by inhibiting the production of extracellular polysaccharides (EPS), the key components of the biofilm matrix. The *brpA* encodes the surface-associated biofilm-regulatory protein that is vital to biofilm formation and the responses to the environmental stresses (Wen and Burne, 2004). The LytR-CpsA-Psr family proteins encoded by *brpA*. LytR has an important role in the cell division and in sucrose-independent attachment (Chatfield et al., 2005; Yoshida and Kuramitsu, 2002). The decreased expression of *brpA* gene by PSV compromised the stress resistance of *S. mutans*. The *vicR* gene encodes a two-component regulatory system that regulates the expression of several genes related to the sucrose-dependent adherence and the synthesis of polysaccharides, including *ftf* and *gtf* gene family (Senadheera et al., 2005). The *vicR* gene is down-regulated by PSV and the suppression of this gene led to an anti-biofilm forming effect through the inhibition of adherence activity. Various factors, which include environmental stresses and intracellular processes such as carbohydrate metabolism can influence the expression of GTFs (Banas and Vickerman, 2003). Quorum-sensing is an important mechanism in the formation of biofilms by *S. mutans*. The *relA* and *comDE* genes contribute to the quorum-sensing and biofilm formation by *S. mutans*. The *relA* gene encodes RelA, a carbohydrate phosphotransferase system (PTS) protein that regulates the glucose uptake system by phosphoenolpyruvate (Lemos et al., 2004). The *comDE* gene encodes two-component signal transduction systems composed of comD and comE that are part of competence-stimulating peptides. The comD and comE act as histidine kinase receptors and cognate response-regulators in the quorum-sensing of *S. mutans* (Suntharalingam and Cvitkovitch, 2005). If comD and comE are defective, biofilms with reduced biomass can be formed and initial sucrose-independent attachment (Li et al., 2002). The expression of *relA* and *comDE* were also down-regulated, especially the expression of *comDE*, by PSV and the decrease was over 90% in all strains. These results indicated that PSV inhibited the biofilm-forming activity of *S. mutans* by controlling the regulatory genes involved in the quorum-sensing system and the production of the key components for biofilm matrix formation.

Conclusion

It can be concluded that PSV showed cariostatic properties through the following mechanisms: 1) inhibiting biofilm formation by *S. mutans* and the degradation of pre-formed biofilms, 2) inhibiting the production of EPS, and 3) inhibiting bacterial adhesion to the surfaces, and all these inhibiting activities of PSV are through the control of gene expression in *S. mutans*. PSV down-regulated the expression of genes related to the production of EPS, autoinducers of quorum sensing, and the key components of the biofilm matrix. Thus, PSV has high potentials to be used as a treatment as well as a preventive agent for dental caries.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through High Value-added Food Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA)(118037-3).

Author Contributions

Conceptualization: Ahn DU, Paik HD. Data curation: Kim HJ, Lee JH. Formal analysis: Kim HJ, Lee JH. Methodology: Kim HJ, Lee JH. Validation: Lee JH. Writing - original draft: Kim HJ. Writing - review & editing: Kim HJ, Lee JH, Ahn DU, Paik HD.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Abe Y, Itoh T, Adachi S. 1982. Fractionation and characterization of hens egg yolk phosvitin. *J Food Sci* 47:1903-1907.
- Al-Sohaibani S, Murugan K. 2012. Anti-biofilm activity of *Salvadora persica* on cariogenic isolates of *Streptococcus mutans*: *In vitro* and molecular docking studies. *Biofouling* 28:29-38.
- Banas J. 2004. Virulence properties of *Streptococcus mutans*. *Front Biosci* 9:1267-1277.
- Banas JA, Vickerman MM. 2003. Glucan binding proteins of the oral streptococci. *Crit Rev Oral Biol Med* 14:89-99.
- Bowen WH, Koo H. 2011. Biology of *Streptococcus mutans*-derived glucosyltransferases: Role in extracellular matrix formation of cariogenic biofilms. *Caries Res* 45:69-86.
- Bradshaw DJ, Lynch RJ. 2013. Diet and the microbial aetiology of dental caries: New paradigms. *Int Dent J* 63:64-72.
- Burne RA, Chen YY, Wexler DL, Kuramitsu H, Bowen WH. 1996. Cariogenicity of *Streptococcus mutans* strains with defects in fructan metabolism assessed in a program-fed specific-pathogen-free rat model. *J Dent Res* 75:1572-1577.
- Burne RA. 1998. Oral streptococci products of their environment. *J Dent Res* 77:445-452.
- Chatfield CH, Koo H, Quivey RG. 2005. The putative autolysin regulator LytR in *Streptococcus mutans* plays a role in cell division and is growth-phase regulated. *Microbiology* 151:625-631.
- Choi IW, Jung CH, Seog HM, Choi HD. 2004. Purification of phosvitin from egg yolk and determination of its physicochemical properties. *J Food Sci Biotechnol* 13:434-437.
- Daglia M, Stauder M, Papetti A, Signoretto C, Giusto G, Canepari P, Gazzani G. 2010. Isolation of red wine components with anti-adhesion and anti-biofilm activity against *Streptococcus mutans*. *Food Chem* 119:1182-1188.
- Flemming HC, Wingender J. 2010. The biofilm matrix. *Nat Rev Microbiol* 8:623-633.
- Gazzani G, Daglia M, Papetti A. 2012. Food components with anticaries activity. *Curr Opin Biotechnol* 23:153-159.
- Giacaman RA, Jobet-Vila P, Muñoz-Sandoval C. 2019. Anticaries activity of egg ovalbumin in an experimental caries biofilm model on enamel and dentin. *Clin Oral Investig* 23:3509-3516.
- Hamada S, Slade HD. 1980. Biology, immunology, and cariogenicity of *Streptococcus mutans*. *Microbiol Rev* 44:331-384.
- Hanada N, Kuramitsu HK. 1988. Isolation and characterization of the *Streptococcus mutans gtfC* gene, coding for synthesis of both soluble and insoluble glucans. *Infect Immun* 56:1999-2005.
- Islam B, Khan SN, Haque I, Alam M, Mushfiq M, Khan AU. 2008. Novel anti-adherence activity of mulberry leaves: Inhibition of *Streptococcus mutans* biofilm by 1-deoxynojirimycin isolated from *Morus alba*. *J Antimicrob Chemother* 62:751-757.
- Koo H, Hayacibara MF, Schobel BD, Cury JA, Rosalen PL, Park YK, Vacca-smith AM, Bowen WH. 2003. Inhibition of

- Streptococcus mutans* biofilm accumulation and polysaccharide production by apigenin and tt-farnesol. J Antimicrob Chemother 52:782-789.
- Koolheat N, Kamuthachad L, Anthapanya M, Samakchan N, Sranujit RP, Potup P. 2016. Kaffir lime leaves extract inhibits biofilm formation by *Streptococcus mutans*. Nutrition 32:486-490.
- Kulshrestha S, Khan S, Hasan S, Khan ME, Misba L, Khan AU. 2016. Calcium fluoride nanoparticles induced suppression of *Streptococcus mutans* biofilm: An *in vitro* and *in vivo* approach. Appl Microbiol Biotechnol 100:1901-1914.
- Lee HY, Abeyrathne ED, Choi I, Suh JW, Ahn DU. 2014. Sequential separation of immunoglobulin Y and phosvitin from chicken egg yolk without using organic solvents. Poult Sci 93:2668-2677.
- Lee SK, Han JH, Decker EA. 2002. Antioxidant activity of phosvitin in phosphatidylcholine liposomes and meat model systems. J Food Sci 67:37-41.
- Lemos JA, Brown TA, Burne RA. 2004. Effects of relA on key virulence properties of planktonic and biofilm populations of *Streptococcus mutans*. Infect Immun 72:1431-1440.
- Li YH, Tang N, Aspiras MB, Lau PCY, Lee JH, Ellen RP, Cvitkovitch DG. 2002. A quorum-sensing signaling system essential for genetic competence in *Streptococcus mutans* is involved in biofilm formation. J Bacteriol 184:2699-2708.
- Mine Y. 2007. Egg proteins and peptides in human health-chemistry, bioactivity and production. Curr Pharm Des 13:875-884.
- Paes Leme AF, Koo H, Bellato CM, Bedi G, Cury JA. 2006. The role of sucrose in cariogenic dental biofilm formation-new insight. J Dent Res 85:878-887.
- Rajaraman S, Subbiahdoss G, Patchirajan P. 2014. Effect of hen egg white on microbial adhesion and biofilm growth of biomaterial associated infection causing pathogens. Int J Bio-Sci Bio-Technol 6:99-106.
- Russell RR, Mukasa H, Shimamura A, Ferretti JJ. 1988. *Streptococcus mutans* *gtfA* gene specifies sucrose phosphorylase. Infect Immun 56:2763-2765.
- Samaraweera H, Zhang W, Lee EJ, Ahn DU. 2011. Egg yolk phosvitin and functional phosphopeptides: Review. J Food Sci 76:R143-R150.
- Senadheera MD, Guggenheim B, Spatafora GA, Huang YC, Choi J, Hung DC, Treglown JS, Goodman SD, Ellen RP, Cvitkovitch DG. 2005. A VicRK, signal transduction system in *Streptococcus mutans* affects *gtfBCD*, *gbpB*, and *fjt* expression, biofilm formation, and genetic competence development. J Bacteriol 187:4064-4076.
- Shapiro JA. 1998. Thinking about bacterial populations as multicellular organisms. Annu Rev Microbiol 52:81-104.
- Shemesh M, Steinberg D. 2006. Surface plasmon resonance for real-time evaluation of immobilized fructosyltransferase activity. J Microbiol Methods 64:411-415.
- Siriwardhana N, Kalupahana NS, Cekanova M, LeMieux M, Greer B, Moustaid-Moussa N. 2013. Modulation of adipose tissue inflammation by bioactive food compounds. J Nutr Biochem 24:613-623.
- Smalley JW, Birss AJ, McKee AS, Marsh PD. 1995. Congo red binding by *Porphyromonas gingivalis* is mediated by a 66 kDa outer-membrane protein. Microbiology 141:205-211.
- Smith DJ, Anderson JM, King WF, van Houte J, Taubman MA. 1993. Oral streptococcal colonization of infants. Oral Microbiol Immunol 8:1-4.
- Staat RH, Langley SD, Doyle RJ. 1980. *Streptococcus mutans* adherence: Presumptive evidence for protein-mediated attachment followed by glucan-dependent cellular accumulation. Infect Immun 27:675-681.
- Stewart PS. 1996. Theoretical aspects of antibiotic diffusion into microbial biofilm. Antimicrob Agents Chemother 40:2517-2522.

- Suntharalingam P, Cvitkovitch DG. 2005. Quorum sensing in streptococcal biofilm formation. *Trends Microbiol* 13:3-6.
- Sutherland IW. 2001. The biofilm matrix: An immobilized but dynamic microbial environment. *Trends Microbiol* 9:222-227.
- Taborsky G, Mok C. 1967. Phosvitin homogeneity and molecular weight. *J Biol Chem* 242:1495-1501.
- Wen ZT, Burne RA. 2004. LuxS-mediated signaling in *Streptococcus mutans* is involved in regulation of acid and oxidative stress tolerance and biofilm formation. *J Bacteriol* 186:2682-2691.
- Wen ZT, Yates D, Ahn SJ, Burne RA. 2010. Biofilm formation and virulence expression by *Streptococcus mutans* are altered when grown in dual-species model. *BMC Microbiol* 10:111.
- Yoshida A, Kuramitsu HK. 2002. Multiple *Streptococcus mutans* genes are involved in biofilm formation. *Appl Environ Microbiol* 68:6283-6291.